



Sea Surface Height (SSH)
Northbound



Sea Surface Height (SSH)
Southbound



Internal Tide Refraction and Attenuation in the North Pacific

Matthew Alford

Scripps Institution of Oceanography

Zhongxiang Zhao, Luc Rainville, James Girtton

Applied Physics Laboratory / University of Washington

Harper Simmons

University of Alaska / Fairbanks


OSTST meeting, KONSTANZ, October 2014

REFERENCES

Altimetry data provided by Z. Zhao

Model cross section adapted from - J. Klymak


Image design and creation: M. Alford and Center for Environmental Visualization (CEV)



The ocean beneath


SSHA: 25 cm
peak-peak!

isopycnals: 500-
m peak-peak



Internal Waves in Straits Experiment

Alford et al, JPO 2010



Breaking internal lee waves at Kaena Ridge

Pickering et al, in prep JPO 2013

INTERNAL TIDES ESTIMATED FROM SATELLITE ALTIMETRY

WAVECHASERS

TRACKING INTERNAL WAVES AND TURBULENCE ACROSS THE GLOBE

Sea Surface Height (SSH)
Northbound



Sea Surface Height (SSH)
Southbound




Tidal beams from the
Hawaiian & Aleutian ridges

REFERENCES

Altimetry data provided by Z. Zhao

Model cross section adapted from - J. Klymak

Visual design and creation: M. Alford and Center for Environmental Visualization (CEV)



Back


CEV



wavechasers.uw.edu

WAVECHASERS

TRACKING INTERNAL WAVES AND TURBULENCE ACROSS THE GLOBE




• SHOALING INTERNAL TIDES MIX STRONGLY ON CONTINENTAL SLOPES

REFERENCES

Altimetry data provided by Z. Zhao

Model cross section adapted from - J. Klymak

Visual design and creation: M. Alford and Center for Environmental Visualization (CEV)




CEV




wavechasers.uw.edu

Sea Surface Height (SSH)
Northbound




Sea Surface Height (SSH)
Southbound




Distribution of Mixing in models matters.

REFERENCES
Altimetry data provided by Z. Zhao
Model cross section adapted from - J. Klymak
Image design and creation: M. Alford and Center for Environmental Visualization (CEV)

MOC change with tidal mixing




[Jayne 09, CCSM]




[Hallberg, MOM]

Sea Surface Height (SSH)
Northbound



Sea Surface Height (SSH)
Southbound



What leads to the attenuation?

- Loss of coherence (not real attenuation)
- Cylindrical Spreading
- Real attenuation:
 - Parametric Subharmonic instability (PSI) $|\text{lat}| < 28.8$
 - Scattering
 - Wave-wave interactions
 - Other processes?


REFERENCES

Altimetry data provided by Z. Zhao


Model cross section adapted from - J. Klymak

Image design and creation: M. Alford and Center for Environmental Visualization (CEV)

Sea Surface Height (SSH)
Northbound



Sea Surface Height (SSH)
Southbound



Approach

- Loss of coherence (not real attenuation): **harmonic analysis on a 1-year GCM**
- Cylindrical Spreading: **integrate in wedges**
- Real attenuation: **What's LEFT!**
 - Parametric Subharmonic instability (PSI) $|\text{lat}| < 28.8$
 - Scattering
 - Wave-wave interactions


REFERENCES

Altimetry data provided by Z. Zhao

Model cross section adapted from - J. Klymak


Image design and creation: M. Alford and Center for Environmental Visualization (CEV)

Sea Surface Height (SSH)
Northbound



1/8° GCM with tides (GOLD;
Harper Simmons)

Sea Surface Height (SSH)
Southbound




TOTAL: 2-day
harmonic fits


INCOHERENT
: subtract 365-
day harmonic
fits

REFERENCES
Altimetry data provided by Z. Zhao
Model cross section adapted from - J. Klymak
Image design and creation: M. Alford and Center for Environmental Visualization (CEV)


Total energy fluxes, DT = 2 days




Incoherent total M2 energy fluxes, (2 days) - (364 days)



Sea Surface Height (SSH)
Northbound




Sea Surface Height (SSH)
Southbound



- Aleutians
- Hawaii North
- Hawaii South
- Mendocino North
- Mendocino South

- NO PSI, SMOOTH
- NO PSI > 500 KM, SMOOTH
- PSI, ROUGH
- NO PSI, ROUGH
- NO PSI, SMOOTH

Altimetry



Model


REFERENCES

Altimetry data provided by Z. Zhao


Model cross section adapted from - J. Klymak

Image design and creation: M. Alford and Center for Environmental Visualization (CEV)

Sea Surface Height (SSH)
Northbound



Sea Surface Height (SSH)
Southbound




Separate North and South


Altimetry:
total




REFERENCES
Altimetry data provided by Z. Zhao
Model cross section adapted from - J. Klymak
Image design and creation: M. Alford and Center for Environmental Vis



Sea Surface Height (SSH)
Northbound



Sea Surface Height (SSH)
Southbound



Separate North and South

Altimetry:
northward




REFERENCES


Altimetry data provided by Z. Zhao

Model cross section adapted from - J. Klymak


Image design and creation: M. Alford and Center for Environmental Vi



Sea Surface Height (SSH)
Northbound



Sea Surface Height (SSH)
Southbound



Separate North and South

Altimetry:
southward


Depth (m)
0
1000
2000
3000
4000

REFERENCES

Altimetry data provided by Z. Zhao


Model cross section adapted from - J. Klymak

Image design and creation: M. Alford and Center for Environmental V





Coherence


Sea Surface Height (SSH)
Northbound



Sea Surface Height (SSH)
Southbound




Mesoscale rms and mean currents




REFERENCES
 Altimetry data provided by Z. Zhao
 Model cross section adapted from - J. Klymak
 Image design and creation: M. Alford and Center for Environmental Visualization

Sea Surface Height (SSH)
Northbound



Sea Surface Height (SSH)
Southbound




- Aleutians
- Hawaii North
- Hawaii South
- Mendocino North
- Mendocino South


- NO PSI, SMOOTH
- NO PSI > 500 KM, SMOOTH
- PSI, ROUGH
- NO PSI, ROUGH
- NO PSI, SMOOTH

REFERENCES
Altimetry data provided by Z. Zhao
Model cross section adapted from - J. Klymak
Image design and creation: M. Alford and Center for Environmental Visualization (CEV)


Final result: attenuation



Sea Surface Height (SSH)
Northbound



Sea Surface Height (SSH)
Southbound



Conclusions


- INTERNAL TIDE IN THE NORTH PACIFIC ESTIMATED FROM ALTIMETRY
- COHERENCE COMPUTED FROM A 1-YEAR OUTPUT OF A GCM (GOLD) VARIES FROM 0.5 IN STRONGLY EDDYING REGIONS TO >0.8 IN QUIET REGIONS
- INTERNAL TIDES PROPAGATE **FAR** IN THE NORTH PACIFIC
- ATTENUATION STRONGEST OVER ROUGH BATHYMETRY AND WHEN PARAMETRIC SUBHARMONIC INSTABILITY (PSI) POSSIBLE.

REFERENCES

Altimetry data provided by Z. Zhao

Model cross section adapted from - J. Klymak

Image design and creation: M. Alford and Center for Environmental Visualization (CEV)



WAVECHASERS

TRACKING INTERNAL WAVES AND TURBULENCE ACROSS THE GLOBE

• FREE AND STANDING WAVES





CEV



wavechasers.uw.edu

WAVECHASERS

TRACKING INTERNAL WAVES AND TURBULENCE ACROSS THE GLOBE



REFERENCES

Altimetry data provided by Z. Zhao


Model cross section adapted from - J. Klymak

Visual design and creation: M. Alford and Center for Environmental Visualization (CEV)




wavechasers.uw.edu

Sea Surface Height (SSH)
Northward




Ferrari and Wunsch 2009




Near-inertial and tidal forcing and lateral energy flux

Alford, Nature 2003




Diffusivity and dissipation rate inferred from Argo vertical strain using a finescale parameterization



WAVECHASERS

TRACKING INTERNAL WAVES AND TURBULENCE ACROSS THE GLOBE

INTERNAL WAVE BEAMS



Tidal beams from the
Hawaiian & Aleutian ridges

REFERENCES

Altimetry data provided by Z. Zhao

Model cross section adapted from - J. Klymak

Visual design and creation: M. Alford and Center for Environmental Visualization (CEV)



wavechasers.uw.edu